

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/042953

International filing date: 22 December 2004 (22.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/532,363
Filing date: 24 December 2003 (24.12.2003)

Date of receipt at the International Bureau: 31 March 2005 (31.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1298977

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

March 21, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/532,363

FILING DATE: *December 24, 2003*

RELATED PCT APPLICATION NUMBER: *PCT/US04/42953*



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office



18379 U.S. PTO

PTO/SB/16 (08-03)

Approved for use through 07/31/2006. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EL 837 182 453

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Michael R.		Rosen		25 East 86th Street New York, New York 10028	
Additional inventors are being named on the _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input type="checkbox"/> Customer Number: _____					
OR					
<input checked="" type="checkbox"/> Firm or Individual Name		John P. White, Esq.			
Address		Cooper & Dunham LLP			
Address		1185 Avenue of the Americas			
City		New York		State	NY
Country		USA		Zip	10036
		Telephone	(212) 278-0400	Fax	(212) 391-0525
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages 14		<input type="checkbox"/> CD(s), Number _____			
<input checked="" type="checkbox"/> Drawing(s) Number of Sheets 4		<input checked="" type="checkbox"/> Other (specify) claims pages 2			
<input type="checkbox"/> Application Date Sheet. See 37 CFR 1.76		Abstract 1 page			
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.		FILING FEE Amount (\$)			
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.		\$80.			
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 03-3125					
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.					
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

[Page 1 of 2]

Respectfully submitted,

SIGNATURE

TYPED or PRINTED NAME Peter J. Phillips

TELEPHONE (212) 278-0400

Date December 24, 2003

REGISTRATION NO. 29,691

(if appropriate)

Docket Number: 0575-71126-Pro

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

331431 U.S. PTO
60/532863

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (08-03)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Docket Number 0575-71126-Pro

INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle [if any])	Family or Surname	Residence (City and either State or Foreign Country)
Peter R.	Brink	101 Dyke Road Setauket, New York 11733
Richard B.	Robinson	67 Roosevelt Street Cresskill, New Jersey 07626
Ira S.	Cohen	23 Hawks Nest Road Stony Brook, New York 11790

[Page 2 of 2]

Number 2 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Michael R. Rosen, Peter R. Brink, Ira S.
Cohen and Richard B. Robinson

Serial No. : Not yet assigned

Filed : December 24, 2003

For : **CREATION OF A BIOLOGICAL ATRIOVENTRICULAR
BYPASS TO COMPENSATE FOR ATRIOVENTRICULAR
BLOCK**


1185 Avenue Of The Americas
New York, New York 10036
December 24, 2003


Mail Stop Provisional Patent Application
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

**EXPRESS MAIL
CERTIFICATE OF MAILING
FOR ABOVE-IDENTIFIED APPLICATION**

"Express Mail" mailing label number: EL 837 182 453 US.
Date of Deposit: December 24, 2003

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 on the date indicated above and is addressed to the Mail Stop Provisional Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.


Printed Name: Trinidad Iscoa


John P. White
Registration No. 28,678
Peter J. Phillips
Registration No. 29,691
Attorney for Applicant
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, NY 10036
(212) 278-0428

Application
for
United States Letters Patent

To all whom it may concern

Be it known that

MICHAEL R. ROSEN PETER R. BRINK; IRA S. COHEN;
RICHARD B. ROBINSON;

have invented certain new and useful improvements in

CREATION OF A BIOLOGICAL ATRIOVENTRICULAR BYPASS TO
COMPENSATE FOR ATRIOVENTRICULAR BLOCK

of which the following is a full, clear and exact description

5 **CREATION OF A BIOLOGICAL ATRIOVENTRICULAR BYPASS TO COMPENSATE
FOR ATRIOVENTRICULAR BLOCK**

10 **Statement of Federally Sponsored Research or Development**

Work on this invention was sponsored by USPH5 and NHLBI under
award number HL-28958.

15 **Background of the Invention**

Throughout this application, various publications are
referenced to as footnotes or within parentheses. Disclosures
20 of these publications in their entireties are hereby
incorporated by reference into this application to more fully
describe the state of the art to which this invention
pertains. Full bibliographic citations for these references
may be found at the end of this application, preceding the
25 claims.

One of the major indications for electronic pacemaker therapy
is high degree heart block, such that a normally functioning
sinus node impulse cannot propagate to the ventricle. The
30 result is ventricular arrest and/or fibrillation, and death.

Acute myocardial infarction (MI) afflicts millions of people
each year inducing significant mortality and, in a large number
of survivors, marked reductions in myocyte number and in
35 cardiac pump function. Adult cardiac myocytes divide only
rarely, and the usual response to myocyte cell loss is
hypertrophy that often progresses to congestive heart failure,
a disease with a significant annual mortality. There have been
recent reports of the delivery of mesenchymal stem cells
40 (MSCs a multipotent cell population of blood lineage) to the

5 hearts of post-Mi patients resulting in improved mechanical performance^{1,2}. The presumption in these and other animal studies³, is that the MSCs integrate into the cardiac syncytium and then differentiate into new heart cells restoring mechanical function.

10

5 Summary of the Invention

The present invention uses biological means for cell therapy to build a bypass tract in the heart that will take over the function of a diseased atrioventricular node. Adult human
10 mesenchymal stem cells (hMSCs) may be prepared in one of four ways (see below) and grown in culture on a non-bioreactive material. Once growth is complete the material has one end sutured to the atrium, and the other to the ventricle. Electrical signals generated by the sinus node to activate the
15 atria will propagate across the artificially constructed tract to excite the ventricle as well. In this way the normal sequence of atrioventricular activation will be maintained.

Four methods that may be used for preparing the hMSCs are:

20

1: In culture without incorporation of additional molecular determinants of conduction. Here the cells' own characteristic to generate gap junctions that communicate electrical signals are used as a means to propagate an
25 electronic wave from atrium to ventricle.

30

2: In culture following electroporation to add the gene for connexins 43, 40 and/or 45, the culture's electrotonic propagation of atrial signals to the ventricle.

35

3: In culture following electroporation to add the alpha and the accessory subunits of the L-type calcium channel, thereby increasing the likelihood of not just electrotonic propagation of a wavefront, but its active
propagation by an action potential.

4: A combination of 2 and 3.

5 The preparation of a bypass in this fashion not only will
facilitate propagation from atrium to ventricle, but will
provide sufficient delay from atrial to ventricular
contraction to maximize ventricular filling and emptying. The
goal is to mimic the normal activation and contractile
10 sequence of the heart. Moreover, this approach, when used
with gene therapy and stem cell technology to improve atrial
impulse initiation in the setting of sinus node disease offers
a completely physiologic system rather than its electronic
replacement.

15

According to the invention, a method of creating an
atrioventricular bypass tract for a heart is provided,
comprising growing mesenchymal stem cells into a strip with
two ends, attaching one end of the strip onto the atrium of
20 the heart, and attaching the other end of the strip to the
ventricle of the heart, to create a tract connecting the
atrium to the ventricle to provide a path for electrical
signals generated by the sinus node to propagate across the
tract and excite the ventricle.

25

5 Description of the Drawings

- Figure 1.** Identification of connexins in gap junctions of hMSCs. Immunostaining of Cx43 (A), Cx40 (B) and Cx45 (C). D, Immunoblot analysis of Cx43 in canine ventricle myocytes and hMSCs. Whole cell lysates (120 jig) from ventricle cells or hMSCs were resolved by SDS, transferred to membranes, and blotted with Cx43 antibodies. Migration of molecular weight markers is indicated to the right to the blot.
- Figure 2.** Macroscopic and single channel properties of gap junctions between hMSC pairs. Gap junction currents (I_j) elicited from hMSCs using symmetrical bipolar pulse protocol showed two types of voltage dependent current deactivation: (A)-symmetrical, (B): asymmetrical.
- C,D** Single channel recordings from pairs of hMSCs. Pulse protocol (V_1 and V_2) and associated multichannel currents (I_z) recorded from a cell pair during maintained V_j of ± 80 mV. The discrete current steps indicate the opening and closing of single channels. Dashed line: zero current level. The all points current histograms on the right-hand side revealed a conductance of ~ 50 pS. Glass coverslips with adherent cells were transferred to an experimental chamber perfused at room temperature ($\sim 22^\circ\text{C}$) with bath solution containing (mM): NaCl, 150; KCl, 10; CaCl_2 , 2; HEPES, 5 (pH 7.4); glucose, 5. The patch pipettes were filled with solution containing (mM): K^+ aspartate $^-$, 120; NaCl, 10; MgATP, 3; HEPES, 5 (pH 7.2); EGTA, 10 (pCa ~ 8); filtered through $0.22\ \mu\text{m}$ pores. When filled, the resistance of the pipettes measured 1-2 M Ω . Experiments were carried out on cell pairs using a double voltage-clamp. This method permitted to control the membrane potential (V_m) and measure the associated junctional

5 currents (I_j).

Figure 3. Macroscopic properties of junctions in cell pairs between a hMSC and HeLa cell expressing only Cx40, Cx43 or Cx45. In all cases hMSC to HeLa cell coupling was tested 6 to 12 after hours initiating co-culture.

10 **A,** I_j elicited in response to a series of voltage steps (V_j) in hMSC-HeLaCx43 pairs.

Top: symmetrical current deactivation; bottom: asymmetrical current voltage dependence.

15

B, Macroscopic I_j recordings from hMSC-HeLaCx40 pairs exhibit symmetrical (top panel) and asymmetrical (bottom panel) voltage dependent deactivation.

20 **C,** Asymmetric I_j from hMSC-HeLaCx43 pair exhibits voltage dependent gating when Cx45 side is relative negative. I_j recorded from hMSC.

D, Cell-to-cell LY spread in cell pairs: from a HeLa Cx43 to an hMSC (top panel) and from an hMSC to a HeLa Cx43 to (bottom panel). In both cases a pipette containing 2 mM LY was attached to the left-handed cell in the whole-cell configuration.

30 Epifluorescent micrographs taken at 12 min after dye injection show LY spread to the adjacent (right-handed) cell. The simultaneously measured junctional conductance⁶ revealed g_j of ~16 nS and ~18 nS of the pairs, respectively. Cell Tracker green was used to distinguish
35 hMSCs from HeLa cells or vice versa in all experiments⁸.

Figure 4. Macroscopic and single channel properties of gap junctions between hMSC-canine ventricle cell pairs. Myocytes were plated between 12 and 72 hours and co-cultured with hMSCs for 6 to 12 hours before measuring coupling. **A**, Top panel: Phase-contrast micrograph of a hMSC-canine ventricle pair. Bottom pane: Monopolar pulse protocol (V_1 and V_2) and associated macroscopic junctional currents (I_z) exhibiting asymmetrical voltage dependence. **B**, Top panel: Multichannel current elicited by symmetrical biphasic 60 mV pulse. Dashed line, zero current level; dotted lines, represent discrete current steps indicative of opening and closing of channels. The current histograms yielded a conductance of ~ 40 -50 pS. Bottom panel: Multichannel recording during maintained V_j of 60 mV. The current histograms revealed several conductances of 48 to 64 pS with several events with conductance of 84 pS to with 99 pS (arrows) which resemble operation of Cx43, heterotypic Cx40-Cx43 and/ or homotypic Cx40 channels.

5 Description of the Invention

According to the invention, a method of creating an atrioventricular bypass tract for a heart is provided, comprising growing mesenchymal stem cells into a strip with
10 two ends, attaching one end of the strip onto the atrium of the heart, and attaching the other end of the strip to the ventricle of the heart, to create a tract connecting the atrium to the ventricle to provide a path for electrical signals generated by the sinus node to propagate across the
15 tract and excite the ventricle.

The steps of attaching may be performed by suturing. The stem cells may be adult human mesenchymal stem cells. The step of growing may comprise growing the stem cells in culture on a
20 non-bioreactive material. The step of growing may be performed in an environment substantially free of any additional molecular determinants of conduction.

The method may further comprise a step of adding a gene to the
25 mesenchymal stem cells by electroporation. The gene may encode for a connexin, such as connexin 40, connexin 43, and/or connexin 45. The step of adding a gene by electroporation may include adding alpha and accessory subunits of L-type calcium. The step of adding a gene by
30 electroporation may include adding the gene for connexions and adding alpha and accessory subunits of L-type calcium channel.

MSCs express connexins that are the building block proteins of
35 gap junctions and can form functional gap junctions with one another, with cell lines expressing cardiac connexins, and with adult cardiac myocytes. Further, the connexins expressed

5 suggest that hMSCs should readily integrate into electrical syncytia of many tissues promoting repair or serving as the substrate for a therapeutic delivery system.

Human mesenchymal stem cells (Poietics™ hMSCs - Mesenchymal
10 stem cells, Human Bone Marrow) were purchased from Clonetics/BioWhittaker (Walkersville, M.D.) and cultured in MCS growing media and used from passages 2-4. Typical punctate staining for Cx43 and Cx40 was seen along regions of intimate cell to cell contact of the MSCs grown in culture as
15 monolayers (Figure 1 A,B). Cx45 staining was also detected but unlike that of Cx43 or Cx40 was not typical of connexin distribution in cells. Rather it was characterized by fine granular cytoplasmic and reticular-like staining with no readily observed membrane associated plaques (Figure 1C).
20 This does not exclude the possibility that Cx45 channels exist but does imply that their number relative to Cx43 and Cx40 homotypic, heterotypic and heteromeric channels is low. Figure 1D illustrates Western blot analysis⁴ for canine ventricle myocytes and hMSCs with a Cx43 polyclonal antibody
25 which adds further proof of Cx43 presence in hMSCs.

Gap junctional coupling among hMSCs is demonstrated in Figure 2. Junctional currents recorded between hMSC pairs show quasi-symmetrical (Figure 2A) and asymmetrical (Figure 2B) voltage
30 dependency arising in response to symmetric transjunctional voltage steps of equal amplitude but opposite sign. These behaviors are typically observed in cells which co-express Cx43 and Cx40⁴.

35 Figures 2C and 2D illustrate typical multichannel recordings from a hMSC pair. Using 120 mM K aspartate as a pipette solution channels were observed with unitary conductances of

5 28-80 pS range. Operation of channels with ~50 pS conductance (see Figure 2 C) is consistent with previously published values^{5,6} for Cx43 homotypic channels. This does not preclude the presence of other channel types, it merely suggests that Cx43 forms functional channels in hMSCs.

10

To further define the nature of the coupling hMSCs were co-cultured with human HeLa cells stably transfected with Cx43, Cx40, and Cx45⁷ and it was found that hMSCs were able to couple to all these transfectants. Figure 3A shows an example of
15 junctional currents recorded between an hMSC and HeLaCx43 cell pairs that manifested symmetrically and asymmetrically voltage dependent currents. The quasi-symmetric record suggests that the dominant functional channel is homotypic Cx43 while the asymmetric record suggests the activity of
20 another connexin in the hMSC (presumably Cx40 as shown by immunohistochemistry, see Figure 1) that could be either a heterotypic or heteromeric form or both. These records are similar to those published for transfected cells: heterotypic and mixed (heteromeric) forms of Cx40 and Cx43^{4,8}.
25 Co-culture of hMSCs with HeLa cells transfected with Cx40 (Figure 3B) also revealed symmetric and asymmetric voltage dependent junctional currents consistent with the co-expression of Cx43 and Cx40 in the hMSCs similar to the data for Cx43 HeLa-hMSC pairs. HeLa cells transfected with
30 Cx45 coupled to hMSCs always produced asymmetric junctional currents with pronounced voltage gating when Cx45 (HeLa) side was negative (Figure 3C). This is consistent with the dominant channel forms in the hMSC being Cx43 and Cx40 as both produce asymmetric currents when they form heterotypic
35 channels with Cx45^{4,8}. This does not exclude Cx45 as a functioning channel in hMSCs but it does indicate that Cx45 is a minor contributor to cell to cell coupling in hMSCs.

5 The lack of visualized plaques in the immunostaining for Cx45 (Figure 1) further supports this interpretation.

Figure 3D shows Lucifer Yellow transfer from HeLaCx43 cell to an hMSC cell (top panel) and transfer from an hMSC to a HeLaCx43 (bottom panel). The junctional conductance of the cell pairs was simultaneously measured by methods described earlier⁶ and revealed conductances of ~ 16 nS and ~18 nS, respectively. The transfer of Lucifer Yellow was similar to that previously reported for homotypic Cx43 or co-expressed Cx43 and Cx40 in HeLa cells⁶. Cell Tracker green (Molecular Probes) was always used in one of the two populations of cells to allow heterologous pairs to be identified⁸.

hMSCs were also co-cultured with adult canine ventricular myocytes. As shown in Figure 4 the hMSCs couple electrically with cardiac myocytes. Both macroscopic (Figure 4A) and multichannel (Figure 4B) records were obtained. Junctional currents in Figure 4A are asymmetric while those in Figure 4B show unitary events of the size range typically resulting from the operation of homotypic Cx43 or heterotypic Cx43-Cx40 or homotypic Cx40 channels^{4,8}. Heteromeric forms are also possible whose conductances are the same or similar to homotypic or heterotypic forms.

30 In studies of cell pairs were demonstrated effective coupling of hMSC to other hMSC (13.8 ± 2.4 nS, $n=14$), to HeLa Cx43 (7.9 ± 2.1 nS, $n=7$), to HeLa Cx40 (4.6 ± 2.6 nS, $n=5$), to HeLa Cx45 (11 ± 2.6 nS, $n=5$) and to ventricular myocytes (1.5 ± 1.3 nS, $n=4$). Results show that hMSCs couple to one another via Cx43 and Cx40. In addition, they form functional gap junction channels with cells transfected with Cx43, Cx40 or Cx45 as well as canine ventricular cardiomyocytes. These

5 data support the possibility of using MSCs as a therapeutic substrate for repair of cardiac tissue. Other syncytia such as vascular smooth muscle or endothelial cells should also be able to couple to the hMSCs because of the ubiquity of Cx43 and Cx40^{9,10}. Thus they may also be amenable to hMSCs
10 based therapeutics, as follows: hMSCs can be transfected to express ion channels which then can influence the surrounding syncytial tissue.

Alternatively, the hMSCs can be transfected to express genes
15 that produce small therapeutic molecules capable of permeating gap junctions and influencing recipient cells. Further, for short term therapy, the small molecules can be directly loaded into hMSCs for delivery to recipient cells. The success of such an approach is dependent on gap junction
20 channels as the final conduit for delivery of the therapeutic agent to the recipient cells. The feasibility of one such approach was demonstrated by transfecting hMSCs with mHCN2, a gene encoding the cardiac pacemaker channel, and delivering them to the canine heart where they generate
25 a spontaneous rhythm.

5 References

1. Strauer, B.E. et al Repair of infarcted myocardium by autologous intracoronary mononuclear bone marrow cell transplantation in humans. *Circulation* 106, 1913-1918 (2002).
- 10 2. Perin, E.C., Geng, Y.J. & Willerson, J.T. Adult stem cell therapy in perspective. *Circulation* 107, 935-938 (2003).
3. Orlic, D. et al. Bone marrow cells regenerate infarcted myocardium. *Nature* 410, 701-705 (2001).
4. Valiunas, V., Gemel, J., Brink, P.R. & Beyer, E.C. Gap junction channels formed by coexpressed connexin40 and connexin43. *Am. J. Physiol. Heart Circ. Physiol.* 2001. Oct.;281. (4.):H1675. -89. 281, H1675-H1689 (2001).
- 15 5. Valiunas, V., Bukauskas, F.F. & Weingart, R. Conductances and selective permeability of connexin43 gap junction channels examined in neonatal rat heart cells. *Circ. Res.* 80, 708-719 (1997).
- 20 6. Valiunas, V., Beyer, E.C. & Brink, P.R. Cardiac gap junction channels show quantitative differences in selectivity. *Circ. Res.* 91, 104-111 (2002).
- 25 7. Elfgang, C. et al. Specific permeability and selective formation of gap junction channels in connexin-transfected HeLa cells. *J. Cell Biol.* 129, 805-817 (1995).
8. Valiunas, V., Weingart, R. & Brink, P.R. Formation of heterotypic gap junction channels by connexins 40 and 43. *Circ. Res.* 2000. Feb.4;86. (2.):E42.-9.86, E42-E49 (2000).
- 30

9. Wang.H.Z. et al. Intercellular communication in cultured human vascular smooth muscle cells. *Am. J. Physiol Cell Physiol* 281, C75-C88 (2001).
10. Beyer.E.C. Gap junctions. *Int. Rev. Cytol.* 137C, 1-37 1993).

5 We claim:

1. A method of creating an atrioventricular bypass tract for a heart, comprising:

10 growing mesenchymal stem cells into a strip with two ends;

 attaching one end of the strip onto the atrium of the heart, and

15 attaching the other end of the strip to the ventricle of the heart, to create a tract connecting the atrium to the ventricle to provide a path for electrical signals generated by the sinus node to propagate across the tract and excite the ventricle.

20 2. The method of claim 1, wherein the steps of attaching are performed by suturing.

3. The method of claim 1, wherein the stem cells are adult human mesenchymal stem cells.

25 4. The method of claim 3, wherein the step of growing comprises growing the stem cells in culture on a non-bioreactive material.

30 5. The method of claim 4, wherein the step of growing is performed in an environment substantially free of any additional molecular determinants of conduction.

35 6. The method of claim 1, further comprising a step of adding a gene to the mesenchymal stem cells by electroporation.

7. The method of claim 6, wherein the gene encodes for a connexin.

10 8. The method of claim 7, wherein the connexin includes connexin 40.

9. The method of claim 7, wherein the connexin includes connexin 43.

15 10. The method of claim 7, wherein the connexin includes connexin 45.

11. The method of claim 6, wherein the step of adding a
20 gene by electroporation includes adding alpha and accessory subunits of L-type calcium.

12. The method of claim 6, wherein the step of adding a
25 gene by electroporation includes adding the gene for connexions and adding alpha and accessory subunits of L-type calcium channel.

5 **CREATION OF A BIOLOGICAL ATRIOVENTRICULAR BYPASS TO COMPENSATE
FOR ATRIOVENTRICULAR BLOCK**

Abstract of the Disclosure

10 A method of creating an atrioventricular bypass tract for a
heart comprises growing mesenchymal stem cells into a strip
with two ends, attaching one end of the strip onto the atrium
of the heart, and attaching the other end of the strip to the
ventricle of the heart, to create a tract connecting the
atrium to the ventricle to provide a path for electrical
15 signals generated by the sinus node to propagate across the
tract and excite the ventricle.







FIGURE 1







FIGURE 2





FIGURE 3

BEST AVAILABLE COPY


A

hMSC



canine ventricle cell

B



BEST AVAILABLE COPY





FIGURE 4